

Light in honeycomb atomic lattice

8/12/2022

Pierre Wulles



anr[®]

Phd student working with **Sergey Skipetrov**:



In Grenoble at LPMMC:



Overview of the system




Figure 1.



- No nearest-neighbour model
- No crystal (no “bonds” between sites)
- Sites B and magnetic field \mathbf{B} are two unrelated things







Figure 2. **Ultracold atoms in a honeycomb optical lattice**

Example by Mattis Reisner, Matthieu Bellec, Ulrich Kuhl, and Fabrice Mortessagne:



$$\begin{aligned}
 H = & \hbar \sum_{n=1}^N \sum_{\alpha_n = \sigma_{\pm,0}} \left(\omega_0 + \delta\omega_0^{(A,B)} + \text{sgn}(\alpha_n) \mu_B B - i \frac{\Gamma_0}{2} \right) |\alpha_n\rangle \langle \alpha_n| \\
 & + \frac{3\pi\hbar\Gamma_0}{k_0} \sum_{n \neq m}^N \sum_{\alpha_n, \beta_m = \sigma_{\pm,0}} \mathcal{G}_{\alpha\beta}(\mathbf{r}_n - \mathbf{r}_m) |\alpha_n\rangle \langle \beta_m|
 \end{aligned}$$

- N the number of identical two-levels atoms
- $\omega_0 + \delta\omega_0^{(A,B)}$ resonance frequency of atoms of the sublattice A or B
- B the static magnetic field
- $\mathcal{G}_{\alpha\beta}$ the dyadic Green's function describing the coupling of atoms by EM waves
- σ represents the three polarisations \pm in the plane and 0 perpendicular



And we define a $3N \times 3N$ dimensionless matrix G made of 3×3 blocks: $G\Psi = \Lambda\Psi$

We define two very important parameters:

- $\Delta_{AB} = (\delta\omega_0^B - \delta\omega_0^A) / 2\Gamma_0$ Frequency detuning
- $\Delta_{\mathbf{B}} = \mu B / \Gamma_0$ Zeeman Shift

Brillouin Zone




Figure 3. if $k_0 \gtrsim 1/a$ waves can escape

Periodicity + TE modes only:

$$\Psi_n = \psi(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{r}_n}$$

Transforms the problem into:

$$G\Psi = \Lambda\Psi \quad \Rightarrow \quad M(\mathbf{k})\psi(\mathbf{k}) = \Lambda\psi(\mathbf{k})$$

Where M is a 4×4 matrix:

$$M(\mathbf{k}) = \begin{pmatrix} S_1^{\sigma+\sigma+} & S_1^{\sigma+\sigma-} & S_2^{\sigma+\sigma+} & S_2^{\sigma+\sigma-} \\ S_1^{\sigma-\sigma+} & S_1^{\sigma-\sigma-} & S_2^{\sigma-\sigma+} & S_2^{\sigma-\sigma-} \\ S_3^{\sigma+\sigma+} & S_3^{\sigma+\sigma-} & S_4^{\sigma+\sigma+} & S_4^{\sigma+\sigma-} \\ S_3^{\sigma-\sigma+} & S_3^{\sigma-\sigma-} & S_4^{\sigma-\sigma+} & S_4^{\sigma-\sigma-} \end{pmatrix}$$

$$S_1^{\alpha\beta} = \sum_{\mathbf{r}_m \in A} G_{AA}^{\alpha\beta}(\mathbf{r}_m)e^{i\mathbf{k}\cdot\mathbf{r}_m}$$

$$S_2^{\alpha\beta} = \sum_{\mathbf{r}_m \in A} G^{\alpha\beta}(\mathbf{r}_m + \mathbf{a}_1)e^{i\mathbf{k}\cdot\mathbf{r}_m}$$

$$S_3^{\alpha\beta} = \sum_{\mathbf{r}_m \in A} G^{\alpha\beta}(\mathbf{r}_m - \mathbf{a}_1)e^{i\mathbf{k}\cdot\mathbf{r}_m}$$

$$S_4^{\alpha\beta} = \sum_{\mathbf{r}_m \in A} G_{BB}^{\alpha\beta}(\mathbf{r}_m)e^{i\mathbf{k}\cdot\mathbf{r}_m}$$

Using Poisson summation formula:

$$\sum_{\substack{\mathbf{r}_m \in A \\ \mathbf{r}_m \neq 0}} G^{\alpha\beta}(\mathbf{r}_m) e^{i\mathbf{k} \cdot \mathbf{r}_m} = \frac{1}{\mathcal{A}} \sum_{\mathbf{g}_m \in A'} g^{\alpha\beta}(\mathbf{g}_m - \mathbf{k}; 0) - G^{\alpha\beta}(\mathbf{0})$$

To ensure convergence of this integral we introduce a gaussian cut-off :

$$g^{\alpha\beta}(\mathbf{q}; \mathbf{r}') = -\frac{6\pi}{k_0} \frac{(k_0^2 \delta_{\alpha\beta} - q_\alpha q_\beta)}{2\pi k_0^2} e^{i\mathbf{q} \cdot \mathbf{r}'} \int \frac{1}{k_0^2 - \mathbf{q}^2 - q_z^2} dq_z \rightarrow \int \frac{e^{-a_{\text{ho}}^2(\mathbf{q}^2 + q_z^2)/2}}{k_0^2 - \mathbf{q}^2 - q_z^2} dq_z$$

Which can be expressed in a closed form:

$$g^{\alpha\beta}(\mathbf{q}; 0) = -\frac{6\pi}{k_0} (\delta_{\alpha\beta} k_0^2 - q_\alpha q_\beta) \mathcal{I}(\mathbf{q})$$

With:

$$\mathcal{I}(\mathbf{q}) = \chi(\mathbf{q}) \frac{\pi}{\Lambda(\mathbf{q})} (-i + \operatorname{erfi}(a_{\text{ho}} \Lambda(\mathbf{q}) / \sqrt{2}))$$

$$\chi(\mathbf{q}) = \frac{1}{2\pi k_0^2} e^{-(k_0 a_{\text{ho}})^2 / 2}$$

$$\Lambda(\mathbf{q}) = \sqrt{k_0^2 - q^2}$$

And similarly we can compute $G^{\alpha\beta}(\mathbf{0})$ as the inverse Fourier transform of $g^{\alpha\beta}(\mathbf{q}; 0)$ where $\mathbf{q} = \mathbf{0}$

$$G^{\alpha\beta}(\mathbf{0}) = - \left(\frac{\operatorname{erfi}(k_0 a_{\text{ho}} / \sqrt{2}) - i}{e^{k_0^2 a_{\text{ho}}^2 / 2}} - \frac{-1/2 + (k_0 a_{\text{ho}})^2}{\sqrt{\pi/2} (k_0 a_{\text{ho}})^3} \right) \delta_{\alpha\beta}$$

2017, *PRL 119*, J. Perczel, and M. D. Lukin 10.1103/PhysRevLett.119.023603

Only the imaginary part is convergent when $a_{\text{ho}} \rightarrow 0$ and the real part is divergent (Lamb shift):

$$\operatorname{Im}(G_{\alpha\beta}^*(\mathbf{0})) \xrightarrow{a_{\text{ho}} \rightarrow 0} 1$$

Band diagram

By considering eigenfrequencies $\omega = \omega_0 - \frac{\Gamma_0}{2} \Re(\Lambda)$ and decay rates $\Gamma = \Gamma_0 \Im(\Lambda)$ we can plot the band diagram by findings eigenvalues of $M(\mathbf{k})$

Video




Figure 4. $k_0 a = 2\pi \times 0.05$

A compact formula

$$\begin{aligned}
 W_{\text{gap}} &= 2||\Delta_{\mathbf{B}}| - |\Delta_{AB}|| && \text{if } 0 < |\Delta_{\mathbf{B}}| < R_1 \\
 &= \frac{1}{2}(c_0 - c_1 + S - |\Delta_{AB}|) && \text{if } R_1 < |\Delta_{\mathbf{B}}| < R_2 \\
 &= -2|\Delta_{\mathbf{B}}| + S && \text{if } R_2 < |\Delta_{\mathbf{B}}| < R_3 \\
 &= 0 && \text{elsewhere}
 \end{aligned}$$

Where:

$$S = 2\text{Re}\left(\sqrt{4\Delta_{AB}^2 + \frac{1}{4}(c_2 + i c_3)^2}\right)$$


$$R_1 = \frac{1}{4}(c_0 - c_1 + S + 2|\Delta_{AB}|)$$


$$R_2 = \frac{1}{4}(c_0 - c_1 + S - 2|\Delta_{AB}|)$$

$$R_3 = S/2$$

c_0, c_1 and c_2 are elements of the matrix $M(\mathbf{k})$ and they only rely on $k_0 a$, the width of the gap is dependant of three parameters: $\Delta_{\mathbf{B}}$, Δ_{AB} and $k_0 a$.

Numerical confirmation

Figure 5. $k_0 a = 2\pi \times 0.05$





We define the Chern number as:

$$C = \frac{1}{2\pi} \int_{\text{BZ}} \Omega(\mathbf{k}) d^2\mathbf{k}$$

Where $\Omega(\mathbf{k})$ is the Berry curvature such that:

$$\Omega(\mathbf{k}) = i \left[\left\langle \frac{\partial \psi(\mathbf{k})}{\partial k_x} \left| \frac{\partial \psi(\mathbf{k})}{\partial k_y} \right\rangle - \left\langle \frac{\partial \psi(\mathbf{k})}{\partial k_y} \left| \frac{\partial \psi(\mathbf{k})}{\partial k_x} \right\rangle \right]$$

To compute it faster we use the approach of *T. Fukui et al* by discretizing the Brillouin Zone.

We define:

$$C_{\text{gap}} = \sum_{b_i < \text{gap}} c(b_i)$$

Which is the quantity we refer to when speaking about the Chern number of our system.




Figure 6. Chern number for $k_0 a = 2\pi \times 0.05$

Localization of modes

For a mode ψ_n we define the weight of the modes on sites A as:

$$P_A^n(\mathbf{k}) = |\psi_n^{(1)}(\mathbf{k})|^2 + |\psi_n^{(2)}(\mathbf{k})|^2$$




Figure 7.




Figure 8.

It translates as the following boundary conditions:

$$\mathbf{B} \cdot \mathbf{n} = 0$$

$$\mathbf{E} \times \mathbf{n} = 0$$

Method of images:

$$G^{\alpha\beta}(\mathbf{r}_m) \rightarrow \sum_{n=-\infty}^{+\infty} (-1)^n G_{\text{free}}^{\alpha\beta}(\mathbf{r}_m^n)$$

We then get:

$$g^{\alpha\beta}(\mathbf{q}; 0) = -\frac{6\pi}{k_0} (\delta_{\alpha\beta} k^2 - q_\alpha q_\beta) \left(\mathcal{I}(\mathbf{q}) + \frac{i}{\sqrt{k_0^2 - q^2} k_0} \frac{1}{1 + e^{-i\sqrt{k_0^2 - q^2} d}} \right), \sqrt{k_0^2 - q^2} d \neq \pi$$

And:

$$G_{\text{plates}}(\mathbf{0}) = - \left(\frac{\operatorname{erfi}(k_0 a / \sqrt{2}) - i}{e^{k_0^2 a^2 / 2}} - \frac{-1/2 + (k_0 a)^2}{\sqrt{\pi/2} (k_0 a)^3} - 3 \left(\frac{\ln(1 + e^{ik_0 d})}{k_0 d} + i \frac{\operatorname{Li}_2(-e^{ik_0 d})}{(k_0 d)^2} - \frac{\operatorname{Li}_3(-e^{ik_0 d})}{(k_0 d)^3} \right) \right) \delta_{\alpha\beta}$$

The imaginary part is simply the decay rate for one atom between two plates, this was already computed before in **1973**, *OPTICS COMMUNICATIONS*, P. Milloni and P. Knight




Figure 9. $k_0d = 2 - k_0a = 2\pi \times 0.05$




Figure 10.

- Study of band diagram and topological properties of honeycomb atomic lattices
- General formula to describe the width of the gap in the optical spectrum of an honeycomb lattice with a magnetic field and two types of atoms
- Interesting topological properties occur only when $k_0 a \lesssim 0.1 \times 2\pi$
- Generalizing to the case of atomic lattice between reflecting plates




Figure 11. $k_0 d = 11$

We define a $3N \times 3N$ dimensionless matrix G made of 3×3 blocks:

$$G_{mn} = \delta_{mn}(i \pm 2\Delta_{AB} - 2\alpha\Delta_B)\mathbb{1}_{3 \times 3} + (1 - \delta_{mn})d_{\text{eg}}A_{mn}d_{\text{eg}}^\dagger$$

With:

- $\Delta_{AB} = (\delta\omega_0^B - \delta\omega_0^A) / 2\Gamma_0$
- $\Delta_B = \mu B / \Gamma_0$ and α is $-1, 1$, and 0 for the three diagonal elements.
- d_{eg} is a transformation matrix to go from cartesian to polar coordinates

And:

$$A_{mn} = (1 - \delta_{mn}) \frac{3 e^{ik_0 r}}{2 k_0 r} \left(P(i k_0 r) \mathbb{1} + Q(i k_0 r) \frac{\mathbf{r}_{mn} \otimes \mathbf{r}_{mn}}{r_{mn}^2} \right)$$

Where:

- $k_0 = \omega_0 / c$
- $\mathbf{r}_{mn} = \mathbf{r}_m - \mathbf{r}_n$
- $P(x) = 1 - 1/x + 1/x^2$ and $Q(x) = -1 + 3/x - 3/x^2$




Figure 12.

Chern Numbers in Discretized Brillouin Zone: Efficient Method of Computing (Spin) Hall Conductances, J. Phys. Soc. Jpn. 74, *Takahiro Fukui, Yasuhiro Hatsugai and Hiroshi Suzuki*